

## CLAIMS:

1. A magnetic resonance imaging system comprising a reconstruction unit arranged to
  - reconstruct a complex image of complex valued pixels from magnetic resonance signals
  - 5 - compute a distribution of phase values of the complex image
  - apply a phase correction to the complex image to form a corrected magnetic resonance image and
  - control the phase correction on the basis of the distribution of phase values of the complex image.
- 10 2. A magnetic resonance imaging system as claimed in Claim 1, wherein the distribution of phase values of the complex image is represented by a histogram of the phase values of the complex image.
- 15 3. A magnetic resonance imaging method as claimed in Claim 2, wherein the phase correction is controlled on the basis of a test function of the histogram.
4. A magnetic resonance imaging system as claimed in Claim 3 wherein the test function of the histogram discriminates, in particular enhances peaks, peaks in the histogram
- 20 from broader distributions.
5. A magnetic resonance imaging system as claimed in Claim 4, wherein the test function is formed by the histogram power function.
- 25 6. A magnetic resonance imaging system as claimed in Claim 1, wherein the reconstruction unit is arranged to make the phase correction on the basis of a polynomial phase correction, said polynomial being represented by its polynomial coefficients.

7. A magnetic resonance imaging method as claimed in Claim 6, wherein the reconstruction unit is arranged to control the phase correction by adjusting polynomial coefficients of the polynomial phase correction.
- 5 8. A magnetic resonance imaging system as claimed in Claim 3, wherein the reconstruction unit is arranged
- to make the phase correction on the basis of a polynomial phase correction and
  - control the phase correction by adjusting polynomial coefficients of the polynomial phase correction so as to optimise the test function.
- 10 9. A magnetic resonance imaging system as claimed in Claim 8, wherein the polynomial coefficients are adjusted by way of a trial and improve algorithm controlled on the basis of the test function.
- 15 10. A magnetic resonance imaging method wherein
- a complex image of complex valued pixels is reconstructed from magnetic resonance signals
  - a distribution of phase values of the complex image is computed
  - a phase correction is applied to the complex image to form a corrected magnetic

20 resonance image and

  - the phase correction is controlled on the basis of the distribution of phase values of the complex image.
11. A computer programme comprising instructions to
- 25
- compute a distribution of phase values of a complex image
  - apply a phase correction to the complex image to form a corrected magnetic resonance image and
  - control the phase correction on the basis of the distribution of phase values of the complex image.